

UNITED STATES PLANT PATENT APPLICATION

of

L. PERNILLE AND MOGENS N. OLESEN

for

MINIATURE ROSE PLANT NAMED

'POULhi015'

SUMMARY OF THE INVENTION

BOTANICAL CLASSIFICATION

Rosa hybrida

VARIETY DENOMINATION

5 'POULhi015'

The present invention constitutes a new and distinct variety of miniature rose plant which originated from a controlled crossing between the female parent, an un-named seedling, and the male parent, an un-named seedling. The two
10 parents were crossed in 1993. The resulting seeds were planted in a controlled environment. The new variety is named 'POULhi015'.

The new rose may be distinguished from its seed parent
15 by the following combination of characteristics:

1. The female seed parent has amber yellow flower color, white 'Poulhi015' has a pink flower color.
2. The female seed parent has a taller growth habit than that of 'Poulhi05'.

20 The new variety may be distinguished from its pollen parent, an un-named seedling, by the following combination of characteristics:

1. The pollen parent has yellow flowers while 'Poulhi015' has pink flowers.

2. The pollen parent has a taller growth habit than that of 'Poulhi015'.

The objective of the hybridization of this rose variety for commercial culture was to create a new and
5 distinct variety with unique qualities, such as:

1. Uniform and abundant pink flowers;
2. Vigorous and compact growth;
3. Year-round flowering under glasshouse conditions;
- 10 4. Suitability for production from softwood cuttings in pots;
5. Durable flowers and foliage which make a variety suitable for distribution in the floral industry.

15 This combination of qualities is not present in previously available commercial cultivars of this type, known to the inventors, and distinguish 'POULhi015' from all other varieties of which we are aware.

As part of their rose development program, L. Pernille
20 Olesen and Mogens N. Olesen germinated the seeds from the aforementioned hybridization and conducted evaluations on the resulting seedlings in a controlled environment in Fredensborg, Denmark.

'POULhi015' was selected by the inventors as a single

plant from the progeny of the hybridization in 1993.

Asexual reproduction of 'POULhi015' by cuttings and traditional budding was first done by L. Pernille and Mogens N. Olesen in their nursery in Fredensborg, Denmark in 1994.

5 This initial and other subsequent propagations conducted in controlled environments have demonstrated that the characteristics of 'POULhi015' are true to type and are transmitted from one generation to the next.

10

BRIEF DESCRIPTION OF THE DRAWING

The accompanying color illustration shows as true as is reasonably possible to obtain in color photographs of this type, the typical characteristics of the buds, flowers,
15 leaves, and stems of 'POULhi015'. Specifically illustrated in the drawing:

Fig 1.1; Open flower, and stem showing open
flower and the attachment of leaves,
peduncle;

20

Fig 1.2; Flower bud partially opened;

Fig 1.3; Flower petals, detached;

Fig 1.4; Sepals, receptacle, and pedicel;

Fig 1.5; Juvenile growth;

Fig 1.6; Mature leaves;

Fig 1.7; Bare stems exhibiting thorns.

DETAILED DESCRIPTION OF THE VARIETY

5 The following is a description of 'POULhi015', as
observed in its growth in a field nursery in Jackson County,
Oregon. Observed plants are 3 months of age and were grown
on *Rosa multiflora* understock. Color references are made
using the Royal Horticultural Society (London, England)
10 Colour Chart, 1995, except where common terms of color are
used.

For a comparison, several physical characteristics of
the rose variety 'Poullute', a rose variety from the same
inventors described and illustrated in U.S. Plant Patent No.
15 10,727 issued 30 April, 1997, are compared to 'Poulhi015'
in Chart 1.

CHART 1

	Characteristic	'POULhi015'	'POULLute'
20	General tonality of open flower	Red Group 38A with light intonations of Red Group 52C	Red Group 36A
25	Petalage	35 petals	30 to 35 petals

Characteristic	'POULhi015'	'POULLute'
Coloration of filaments	Yellow Group 12A	Green-Yellow Group 1B
Open flower diameter	45 mm	40 to 50 mm

10

FLOWER AND FLOWER BUD

Blooming habit: Continuous.

Flower bud:

15

Size: Upon opening, 25 mm in length from base of receptacle to end of bud. Bud diameter is 12 to 15 mm.

Bud form: Pointed ovoid.

20

Bud color: As sepals unfold, Red Group 51C; Red Group 51C to 51D at $\frac{1}{4}$ opening.

Sepals:

Upper Surface:

25

Color: Yellow-Green Group 146A.

Surface: Light pubescence.

Lower Surface:

Color: Yellow-Green Group 146B.

Texture: Smooth with stipitate glands.

5 Shape: Sepal apex is cirrhose.
Base is flat at union with receptacle.

Margins: Margins have strong
10 foliaceous appendages on
three of the five sepals.
Stipitate glands present in
medium quantity.

Size: 24 mm (l) x 8 mm (w).

Receptacle:

15 Surface: Slightly pubescent.

Shape: Funnel shaped.

Size: 5 mm (h) x 6 mm (w).

Color: Yellow-Green Group 144B.

Peduncle:

20 Surface: Smooth and slightly
pubescent.

Length: 25 to 30 mm.

Color: Yellow-Green Group 144B.

Strength: Strong.

	Borne:	In small clusters, 1 to 3 flower buds per stem on average.
5	Flower bloom:	
	Fragrance:	Light rose scent.
	Duration:	The blooms have a duration on the plant of approximately 10 to 14 days.
10		Petals fall cleanly away from plant after flowers have fully matured.
	Size:	Average flower diameter is 45 mm when open. Flower depth is 18 mm.
15	Form:	General shape is cupped with a tight center.
	Shape of flower when viewed from the side:	
20		Upon opening, upper part: Flat.
		Upon opening, lower part: Flattened convex.
		Open flower, upper part: Flat.
		Open flower, lower part: Concave.

Petalage: 30 petals under normal conditions
with 4 petaloids.

5 Color:

Upon opening, petals:

Outermost petals:

10 Outer Side: Red Group 51C with
intonations of Red
Group 54B at basal
zone.

15 Inner Side: Red Group 49C with
intonations of Orange
Group 25C at basal
zone.

Innermost petals:

20 Outer Side: Red Group 51C with
intonations of Red
Group 54B at basal
zone.

Inner Side: Red Group 49C with
intonations of Orange
Group 25C at basal
zone.

Upon opening, basal petal spots:

Outermost petals:

Outer Side: Yellow Group 12B.

Inner Side: Yellow Group 12A.

5 Innermost petals:

Outer Side: Yellow Group 12B.

Inner Side: Yellow Group 12A.

After opening, petals:

Outermost petals:

10 Outer Side: Red Group 48D with
intonations of Red
Group 47D.

Inner Side: Red Group 56B.

Innermost petals:

15 Outer Side: Red Group 38A.

Inner Side: Red Group 36A.

After opening, basal petal spots:

Outermost petals:

Outer Side: Yellow Group 12C.

20 Inner Side: Yellow Group 12B.

Innermost petals:

Outer Side: Yellow Group 12B to
12C.

Inner Side: Yellow Group 12B.

5 **General Tonality:** On open flower Red Group 38A with
very light intonations of Red
Group 52C. No change in the
general tonality at the end of
the 7th day. Afterwards, general
tonality is Red Group 52D.

10 **Petals:**

 Petal Reflex: Slightly.

 Petal Margin: Entire with point in center
 of margin.

 Shape: Generally obovate.

 Apex: Rounded.

15 Base: Acute.

 Size: 20 mm (l) x 10 mm (w).

 Thickness: Thin.

 Arrangement: Not formal.

Petaloids:

20 Quantity: 3 to 5.

 Size: 21 mm (l) x 10 mm (w).

 Shape: Base is acute. Apex is
 round.

 Color:

Upper Surface: Red Group 38B.

Lower Surface: Red Group 38C.

Reproductive Organs:

5	Pistils:	
	Length:	5 mm.
	Quantity:	41 (actual count).
	Pollen:	None observed.
	Anthers:	
10	Size:	2 mm.
	Color:	Yellow Group 12A.
	Quantity:	56 (actual count).
	Filaments:	
	Color:	Yellow Group 12A.
15	Length:	5 to 6 mm.
	Stigmas:	Level with the height of the anthers.
	Color:	Greyed-Yellow Group 160D.
	Styles:	
20	Color:	Greyed-Yellow Group 160D.
	Seed formation:	Not observed.

PLANT

5 **Plant growth:** Compact and bushy. When grown as a
pot plant, the average height of the
plant itself is 30 to 35 cm and the
average width is 35 cm.

Stems:

Color:

10 Young wood: Yellow-Green Group 144B.

Older wood: Yellow-Green Group 144B.

Surface:

Young wood: Smooth.

Older wood: Smooth.

15 Thorns:

Incidence: 8 mm per 10 cm of stem.

Size: 6 mm.

Color: Greyed-Red Group 181A.

Shape: Deeply concave.

20

Plant foliage: Normal number of leaflets on normal
leaves in middle of the stem: 5
leaflets.

Compound Leaf size: 80 mm (l) x 45 mm (w).

Color:

Juvenile foliage:

Upper Leaf Surface: Yellow-Green

5

Group 146A.

Lower Leaf Surface: Yellow-Green

Group 148A.

Mature foliage:

Upper Leaf Surface: Yellow-Green

10

Group 146A.

Lower Leaf Surface: Yellow-Green

Group 147B.

Anthocyanin intonation:

Location: Leaf margins and
rachis of juvenile
foliage.

15

Color: Greyed-Red Group 184A.

Plant leaves and leaflets:

20

Stipules:

Size: 20 mm in length.

Shape: Linear with outward
extending apices.

Quantity: 2 per compound leaf.

	Margins:	Finely serrated with stipitate glands.
	Color:	Yellow-Green Group 144A.
5	Petiole:	
	Length:	25 mm.
	Color:	Yellow-Green Group 144B.
10	Underneath:	Thorns and stipitate glands observed.
	Rachis:	
	Size:	35 mm.
	Color:	Yellow-Green Group 144B.
15	Underneath:	Thorns and stipitate glands observed.
	Leaflet:	
	Size:	28 mm (l)x 18 mm(w).
	Edge:	Finely serrated.
20	Shape:	Generally ovate. Cuspidate leaf apex. Leaf base is rounded.
	Texture:	Smooth.
	Arrangement:	Odd pinnate.

Venation: Reticulate.
Glossiness: Glossy.
Thickness: Thick.

5 **Disease resistance:**

Above average resistance to mildew, black spot, and Botrytis under normal growing conditions in Jackson County, Oregon.

10 **Cold hardiness:**

'Poulhi015' has been found to be cold tolerant to USDA cold hardiness zone 6.